Energy levels, radiative rates, and lifetimes for transitions in W XL

Kanti M. Aggarwal and Francis P. Keenan

Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast,
Belfast BT7 1NN, Northern Ireland, UK.

arXiv:1402.1126v1 [physics.atom-ph] 5 Feb 2014

ABSTRACT

Energy levels and radiative rates are reported for transitions in Br-like W XL, calculated with the general-purpose relativistic atomic structure package (GRASP). Configuration interaction (CI) has been included among 46 configurations (generating 4215 levels) over a wide energy range up to 213 Ryd. However, for conciseness results are only listed for the lowest 360 levels (with energies up to $\sim$ 43 Ryd), which mainly belong to the $4s^24p^5$, $4s^24p^44d$, $4s^24p^44f$, $4s4p^6$, $4p^64d$, $4s4p^54d$, $4s^24p^34d^2$, and $4s^24p^34d4f$ configurations, and provided for four types of transitions, i.e. E1, E2, M1, and M2. Comparisons are made with existing (but limited) results. However, to fully assess the accuracy of our data, analogous calculations have been performed with the flexible atomic code (FAC), including even a larger CI than in GRASP. Our energy levels are estimated to be accurate to better than 0.02 Ryd, whereas results for radiative rates (and lifetimes) should be accurate to better than 20% for a majority of the strong transitions.

Received December 13, 2013; accepted February 5, 2014

Running Title: *K. M. Aggarwal and F. P. Keenan / Atomic Data and Nuclear Data Tables xxx (2014) xxx-xxx*

Contents

1. Introduction 00
2. Energy levels 00
3. Radiative rates 00
4. Lifetimes 00
5. Conclusions 00

Acknowledgments 00

References 00

Explanation of Tables

Tables

1. Configurations and levels of W XL. 00
2. Energies (Ryd) for the lowest 360 levels of W XL and their lifetimes (in s). 00
3. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths 00
   (f_{ij}, dimensionless), line strengths S (in atomic unit) for electric dipole (E1), and 00
   A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions in W XL. 00

1. Introduction

Tungsten (W) is an element of active research for studies of fusion plasmas. Since it is an important constituent of tokamak reactor walls, its importance has further increased with the developing ITER project. Due to the high temperatures of fusion plasmas, atomic data (including energy levels and radiative decay rates) are required for many of W ions, mainly to assess and control the radiation loss. Hence there have been several theoretical studies of W ions – see for example, Fournier [1] and Jonauskas et al. [2]. Similarly, there have been laboratory measurements of W emission lines for many of its ions – see for example, Utter et al. [3] and Clementson et al. [4]. These data, including for W XL, have been compiled by NIST (National Institute of Standards and Technology), and are available at their website `http://physics.nist.gov/PhysRefData/ASD/levels_form.html`.

Compared to other W ions there has been less focus on Br-like W XL. In fact, the only calculation available is by Fournier [1], who reported energies for 29 levels of the $4s^24p^5$, $4s^24p^44d$, and $4s4p^6$ configurations, and a selected few of $4s^24p^44f$, $4s^24p^45d$, and $3d^94s^24p^6$. Similarly, his data for radiative rates (A- values) are limited to a few transitions, while for plasma modelling one normally requires a complete set of A- values for *all* transitions.

W XL is a heavy ion (Z = 74) and hence the inclusion of relativistic effects is very important to accurately determine energy levels and subsequently other parameters. However, for this ion configuration interaction (CI) is equally important. For this reason, Fournier [1] included CI among 10 even and 10 odd configurations – see his Table 1. However, he omitted some important configurations, such as $4s^24p^34d^2$, which generate 141 odd parity levels in the 22–42 Ryd energy range – see Table 1 of Aggarwal and Keenan [5]. Our *aim* is to improve upon these earlier calculations and provide a complete set of atomic data among a larger number of levels.

Before we discuss our work, we note that recently S. Aggarwal et al. [6] calculated energy levels, oscillator strengths, radiative rates, and lifetimes for 33 levels of W XL, belonging to the $4s^24p^5$, $4s^24p^44d$, $4s4p^6$, and $3d^94s^24p^6$ configurations. However, their results are unreliable, with discrepancies in A- values and lifetimes with the present calculations of up to two and four orders of magnitude, respectively, as shown by Aggarwal and Keenan [5]. Therefore, we do not discuss their results further.

For the calculations, we have adopted the GRASP (general-purpose relativistic atomic structure package) code, originally developed by Grant et al. [7], and significantly revised by one of its authors (Dr. P. H. Norrington). This code is referred to as GRASP0 and is available at `http://web.am.qub.ac.uk/DARC/`. It is fully relativistic, based on the jj coupling scheme, and includes higher-order relativistic corrections arising from the Breit interaction and QED (quantum electrodynamics) effects. Results obtained from this code are comparable with other subsequent versions, such as GRASP2K [8]–[9].

2. Energy levels

For our calculations we have used the option of *extended average level* (EAL), in which a weighted (proportional to $2j$+1) trace of the Hamiltonian matrix is minimised. This produces a compromise set of orbitals describing closely-lying states with moderate accuracy, and yields results comparable to other options, such as *average level* (AL) – see for example, Aggarwal et al. for several ions of Kr [10] and Xe [11]. Since CI is very important for W XL as already noted, we have performed a series of calculations with increasing amount of CI with up to 42 configurations, namely $4s^24p^5$, $4s^24p^44d$, $4s^24p^44f$, $4s4p^6$, $3d^94s^24p^6$, $4p^64d/4f$, $4s4p^54d/4f$, $4p^34d^2/4f^2/4d4f$, $4s^24p^24d^3$, $4s^24p4d^4$, $4s^24p^24d^24f$, $4s4p^34d^3$, $4p^54d^2$, $3d^94s^24p^54d/4f$, $3p^53d^{10}4s^24p^6$, $3p^53d^{10}4s^24p^54d/4f$, $4s4p^55\ell$, $4p^65\ell$, $4s^24p^45\ell$, and $3d^94s^24p^55\ell$. These generate a total of 4128 levels and are spread over a wide energy range up to 213 Ryd. Many of these configurations have overlapping energy levels, and hence strongly interact and intermix. The levels each configuration generates and their energy ranges are listed in Table 1. Also listed in the table are the configurations included by Fournier [1].

The importance of CI on energy levels and A- values has been clearly demonstrated in our earlier work [5]. Differences of up to 0.4 Ryd with the energy levels of Fournier [1] were noted for several levels – see Table 2 of [5], mainly due to the larger CI included in our calculations. Discrepancies for A- values were larger, up to a factor of three, for several transitions – see Table 3 of [5]. However, in [5] results were only provided for the lowest 31 levels of the $4s^24p^5$, $4s^24p^44d$, and $4s4p^6$ configurations, and two of $3d^94s^24p^6$, which are common with those of [1]. Similarly, A- values were only reported from the lowest two levels of the $4s^24p^5$ ground configuration. Here we list data for a larger number of levels, up to 360, and in addition four more configurations

have been included, apart from the earlier 42 listed above. These are: ($4s^2 4p^4$) 6s, 6p, 6d and 6f, which generate 87 levels in the 81-106 Ryd energy range and overlap with many of the other (1–42) configurations listed in Table 1. However, we note that the effect of including these additional four configurations is *insignificant* on the energy levels and A- values reported in our earlier work [5].

The 46 configurations of W XL listed in Table 1 generate 4215 levels in total, but for conciseness energies are listed in Table 2 for only the lowest 360 levels (with E up to $\sim$ 43 Ryd), which mostly belong to the $4s^2 4p^5$, $4s^2 4p^4 4d$, $4s^2 4p^4 4f$, $4s4p^6$, $4p^6 4d$, $4s4p^5 4d$, $4s^2 4p^3 4d^2$, and $4s^2 4p^3 4d4f$ configurations. Energies for all levels may be obtained electronically on request from one of the authors (KMA: K.Aggarwal@qub.ac.uk).

For ions where CI is very important, it is very difficult to assign a unique label for each level, because many of these mix strongly – see for example Ti VI [12] and Ti VII [13]. In Table A we list the mixing coefficients for 31 levels of the $4s^2 4p^5$, $4s^2 4^4 4d$, and $4s4p^6$ configurations, and except for the lowest two, almost all others are well mixed. Consequently, some of the levels can be easily interchanged, such as 7/60 ($4s^2 4p^4(^1S)4d\ ^2D_{3/2}$ and $4s^2 4p^4(^3P)4d\ ^4F_{3/2}$), 13/26 ($4s^2 4p^4(^1D)4d\ ^2F_{5/2}$ and $4s^2 4p^4(^1D)4d\ ^2D_{5/2}$), and 16/22 ($4s^2 4p^4(^1D)4d\ ^2D_{3/2}$ and $4s^2 4p^4(^3P)4d\ ^2P_{3/2}$). Therefore, the LSJ designations provided in Table 2 are mainly for guidance and should not be taken as definitive. In Table B we provide similar mixing coefficients for the remaining levels, but only up to 65. These are more strongly mixed than those in Table A – see for example 31 and 48, i.e. $4s^2 4p^3(^2P)4d^2(^3F)\ ^4D^o_{1/2}$ and $4s^2 4p^3(^4S)4d^2(^3F)\ ^6F^o_{1/2}$.

Experimental energies for W XL (see the NIST website) are limited to only a few levels for which detailed comparisons have already been made in our earlier work [5]. In the absence of other theoretical results (except of [1]), it is hence difficult to fully assess the accuracy of our energy levels. To make such an assessment we have therefore employed the *Flexible Atomic Code* (FAC) of Gu [14], available from the website `http://sprg.ssl.berkeley.edu/~mfgu/fac/`. This is also a fully relativistic code which provides results for energy levels and A- values comparable to GRASP, as already shown for several other ions, see for example: Aggarwal et al. for Kr [10] and Xe [11] ions, and the more recent work on Ti ions ([12]–[13] and [15]). A further advantage of this code is its high efficiency which means that significantly large calculations can be performed within a reasonably short time. However, because of the strong mixing for several levels of W XL the identification of a level is comparatively more difficult in the FAC calculations than for GRASP.

As with GRASP, we have performed a series of calculations with the FAC code with increasing amount of CI, but focus only on two. These are: (i) FAC1, which includes the same 4215 levels as in GRASP and (ii) FAC2, which also includes all possible combinations of the $4s^2 4p^3 5\ell\ell'$ and $4s4p^4 5\ell\ell'$ configurations, i.e. 11,849 levels in total. Detailed comparisons for 33 levels of the $4s^2 4p^5$, $4s^2 4p^4 4d$, $4s4p^6$, and $3d^9 4s^2 4p^6$ configurations have already been undertaken in our earlier paper [5], and in Table C we provide a similar comparison for the remaining levels, but only up to 65. No measurements or other theoretical results are available for these levels.

For the levels listed in Table C, the energies obtained in our GRASP and FAC1 calculations are comparable (within 0.01 Ryd) and the orderings are also the same. Similarly, the energies in the FAC2 calculations closely agree with those of FAC1, but are higher by $\sim$ 0.01 Ryd for almost all levels. Therefore, the differences between the GRASP and FAC2 energies are less than 0.02 Ryd, and overall there is not much appreciable effect arising from the extra CI included in the FAC2 calculations. Based on this comparison as well as the one shown in Table 2 of [5], we can confidently state that our energy levels listed in Table 2 are accurate to $\sim$ 0.02 Ryd.

3. Radiative rates

The absorption oscillator strength (f_{ij}), a dimensionless quantity, and radiative rate A_{ji} (in s^{-1}) for a transition $i \rightarrow j$ are related by the following expression:

$$f_{ij} = \frac{mc}{8\pi^2 e^2} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} = 1.49 \times 10^{-16} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} \qquad (1)$$

where m and e are the electron mass and charge, respectively, c the velocity of light, λ_{ji} the transition energy/wavelength in Å, and ω_i and ω_j the statistical weights of the lower i and upper j levels, respectively. Similarly, the oscillator strength f_{ij}

(dimensionless) and the line strength S (in atomic unit, 1 a.u. = 6.460×10^{-36} cm^2 esu^2) are related by the following standard equations:

for the electric dipole (E1) transitions:

$$A_{ji} = \frac{2.0261 \times 10^{18}}{\omega_j \lambda_{ji}^3} S \qquad \text{and} \qquad f_{ij} = \frac{303.75}{\lambda_{ji} \omega_i} S, \tag{2}$$

for the magnetic dipole (M1) transitions:

$$A_{ji} = \frac{2.6974 \times 10^{13}}{\omega_j \lambda_{ji}^3} S \qquad \text{and} \qquad f_{ij} = \frac{4.044 \times 10^{-3}}{\lambda_{ji} \omega_i} S, \tag{3}$$

for the electric quadrupole (E2) transitions:

$$A_{ji} = \frac{1.1199 \times 10^{18}}{\omega_j \lambda_{ji}^5} S \qquad \text{and} \qquad f_{ij} = \frac{167.89}{\lambda_{ji}^3 \omega_i} S, \tag{4}$$

and for the magnetic quadrupole (M2) transitions:

$$A_{ji} = \frac{1.4910 \times 10^{13}}{\omega_j \lambda_{ji}^5} S \qquad \text{and} \qquad f_{ij} = \frac{2.236 \times 10^{-3}}{\lambda_{ji}^3 \omega_i} S. \tag{5}$$

In our calculations with the GRASP code the A- and f- values have been determined in both Babushkin and Coulomb gauges, which are equivalent to the length and velocity forms in the more familiar non-relativistic nomenclature. However, in Table 3 we present results in the length form alone, as the velocity form is generally considered to be comparatively less accurate. Included in this table are the transition energies (ΔE$_{ij}$ in Å), radiative rates (A$_{ji}$ in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S in a.u.) for all 16,260 electric dipole (E1) transitions among the lowest 360 levels of W XL. The *indices* used to represent the lower and upper levels of a transition have already been defined in Table 2. Also, in calculating the above parameters we have used the Breit and QED-corrected theoretical energies/wavelengths as listed in Table 2. However, only A-values are included in Table 3 for the 23,975 electric quadrupole (E2), 16,578 magnetic dipole (M1), and 23,501 magnetic quadrupole (M2) transitions. Corresponding results for f- or S- values can be obtained by using Eqs. (1-5).

A detailed comparison among a variety of f- value calculations with both GRASP and FAC has already been made in our earlier paper [5], but only for transitions from the lowest two levels of the $4s^24p^5$ configuration to higher-lying ones. In Table D we provide a similar comparison for transitions from the lowest three levels of the $4s^24p^44d$ configuration. Included in this table are our results for f- values from the GRASP, FAC1 and FAC2 calculations. As noted earlier for energy levels in Table C, the effect of additional CI in FAC2 is small, because for all transitions (strong as well as weak) the f- values are comparable with those in FAC1. Similarly, there is general agreement (within $\sim$ 20%) between the GRASP and FAC results, particularly for those transitions for which the f- values are significant, i.e. ≥ 0.01. For some weak(er) transitions, such as 3–35/47/62, 4–52, and 5–37/40/47, the differences between the two sets of f- values are up to a factor of two. The f- values from FAC are sometimes higher (e.g. 3–47 and 3–59) and in other instances are lower (e.g. 3–35 and 3–62). Such discrepancies for some weak transitions are common among calculations with different codes, and examples may be found in some of our earlier papers, including that on Mg-like ions [16].

Assessing the accuracy of atomic data is not a trivial task [17]. However, a criterion commonly used to determine the reliability of f- (or A-) values is the agreement between their length and velocity forms. Although good agreement between the

two forms is desirable, it is not a necessary condition for accuracy. This is because different sets of configurations may lead to good agreement between the two forms, but entirely different results in magnitude, not only for the weaker inter-combination transitions, but also the allowed ones which are comparatively stable and larger in magnitude. Examples of this are provided in some of our earlier work, such as [18]–[20]. Nevertheless, in Table D we list the velocity/length ratio of the f- values corresponding to our GRASP calculations. For most transitions the two forms agree within $\sim$ 20%, but discrepancies are larger (up to a factor of four) for some, such as 3–34/35/6, 4–34, and 5–32/37/40. Most of these are weak (f $<$ 0.01), but some are not, such as 5–49 and 5–53. Based on the comparisons shown in Table D as well as in Table 3 of our earlier paper [5], we estimate the accuracy of our radiative data to be better than 20%, for a majority of transitions, particularly the stronger ones.

4. Lifetimes

The lifetime τ of a level j is defined as follows:

$$\tau_j = \frac{1}{\sum_i A_{ji}}. \tag{6}$$

In Table 2 we list lifetimes for all 360 levels from our calculations with the GRASP code, which include A- values from all types of transitions, i.e. E1, E2, M1, and M2. Unfortunately, there are no measurements or other theoretical results available with which to compare the lifetimes. However, we hope the data will be useful for future comparisons and further assessment of the accuracy of our radiative rates.

5. Conclusions

In this work, energy levels and lifetimes obtained with the GRASP code have been reported for the lowest 360 levels of W XL, although calculations have been performed for up to 11,849 levels. Radiative rates, oscillator strengths, and line strengths have also been provided for transitions among these levels. To assess the accuracy of our data, similar calculations have also been performed with the FAC code, as no other results are available in the literature. Based on comparisons of calculations with the two independent codes, our energy levels are assessed to be accurate to $\sim$ 0.02 Ryd, whereas the accuracy for other atomic parameters is $\sim$ 20%.

Acknowledgment

KMA is thankful to AWE Aldermaston for financial support.

References

[1] K. Fournier. At. Data Nucl. Data Tables 68 (1998) 1.

[2] V. Jonauskas, G. Gaigalas, S. Kučas. At. Data Nucl. Data Tables 98 (2012) 19.

[3] S.B. Utter, P. Beiersdorfer, E. Träbert. Can. J. Phys. 80 (2002) 1503.

[4] J. Clementson, P. Beiersdorfer, G.V. Brown, M.F. Gu, H. Lundberg, Y. Podpaly, E. Träbert. Can. J. Phys. 89 (2011) 571.

[5] K.M. Aggarwal, F.P. Keenan. Can. J. Phys. 92 (2014) xxx, http://arxiv.org/abs/1310.6620

[6] S. Aggarwal, A.K.S. Jha, M. Mohan. Can. J. Phys. 91 (2013) 394.

[7] I.P. Grant, B.J. McKenzie, P.H. Norrington, D.F. Mayers, N.C. Pyper. Comput. Phys. Commun. 21 (1980) 207.

[8] P. Jönsson, X. He, C.F. Fischer, I.P. Grant. Comput. Phys. Commun. 177 (2007) 597.

[9] P. Jönsson P, G. Gaigalas, J. Bieroń, C.F. Fischer, I.P. Grant. Comput. Phys. Commun. 184 (2013) 2197.

[10] K.M. Aggarwal, F.P. Keenan, K.D. Lawson. At. Data Nucl. Data Tables 94 (2008) 323.

[11] K.M. Aggarwal, F.P. Keenan, K.D. Lawson. At. Data Nucl. Data Tables 96 (2010) 123.

[12] K.M. Aggarwal, F.P. Keenan. Phys. Scr. 88 (2013) 025302.

[13] K.M. Aggarwal, F.P. Keenan. Phys. Scr. 88 (2013) 065304.

[14] M.F. Gu. Can. J. Phys. 86 (2008) 675.

[15] K.M. Aggarwal, F.P. Keenan. Phys. Scr. 88 (2013) 025303.

[16] K.M. Aggarwal, V. Tayal, G.P. Gupta, F.P. Keenan. At. Data Nucl. Data Tables 93 (2007) 615.

[17] K.M. Aggarwal, F.P. Keenan. Fusion Sci. Tech. 63 (2013) 363.

[18] A. Hibbert. Rep. Prog. Phys. 38 (1975) 1217.

[19] P. Oliver, A. Hibbert. AIP Conf. Proc. 1344 (2011) 70.

[20] K.M. Aggarwal. Astrophys. J. Suppl. 118 (1998) 589.

Table A. Mixing coefficients (MC) for the levels of the $4s^24p^5$,$4s4p^6$, and $4s^24p^44d$ configurations of W XL. Numbers outside and inside a bracket correspond to MC and the level, respectively. Level orderings correspond to those of Table 2.

Index	Configuration	Level	Mixing coefficients
1	$4s^24p^5$	$^2P^o_{3/2}$	0.99(1)
2	$4s^24p^5$	$^2P^o_{1/2}$	0.99(2)
3	$4s^24p^4(^3P)4d$	$^4D_{3/2}$	−0.61(3)+0.43(12)−0.29(28)+0.44(16)−0.26(20)−0.23(7)
4	$4s^24p^4(^3P)4d$	$^4P_{1/2}$	0.37(15)−0.68(4)−0.24(9)+0.45(29)−0.32(23)
5	$4s^24p^4(^3P)4d$	$^4D_{5/2}$	−0.35(17) 0.66(5)−0.26(11)+0.26(21)+0.40(13)−0.33(26)
6	$4s^24p^4(^3P)4d$	$^4F_{7/2}$	0.56(6)−0.31(19)+0.52(8)+0.52(18)−0.20(27)
7	$4s^24p^4(^1S)4d$	$^2D_{3/2}$	0.44(60)+0.33(12)+0.23(28)+0.21(22)+0.74(7)
8	$4s^24p^4(^3P)4d$	$^2F_{7/2}$	0.65(19)+0.51(8)+0.25(18)+0.47(27)
9	$4s^24p^4(^3P)4d$	$^2P_{1/2}$	0.25(15)+0.27(4)−0.60(9)+0.36(29)+0.48(23)−0.38(14)
10	$4s^24p^4(^3P)4d$	$^4F_{9/2}$	0.83(10)+0.55(24)
11	$4s^24p^4(^3P)4d$	$^4P_{5/2}$	0.25(17)+0.46(11)−0.39(25)+0.71(66)
12	$4s^24p^4(^3P)4d$	$^4P_{3/2}$	0.26(60)−0.49(12)−0.42(28)+0.42(22)+0.40(16)+0.38(20)
13	$4s^24p^4(^1D)4d$	$^2F_{5/2}$	−0.35(17)−0.39(5)+0.26(11)−0.22(25)+0.48(21)+0.43(13)+0.30(26)−0.32(66)
14	$4s4p^6$	$^2S_{1/2}$	0.45(4)−0.28(9)+0.82(14)
15	$4s^24p^4(^3P)4d$	$^4D_{1/2}$	−0.89(15)−0.31(9)+0.27(29)
16	$4s^24p^4(^1D)4d$	$^2D_{3/2}$	−0.53(60)+0.58(3)+0.21(12)+0.35(22)+0.31(16)−0.30(20)
17	$4s^24p^4(^3P)4d$	$^4F_{5/2}$	0.68(17)+0.35(25)+0.27(21)+0.50(13)−0.26(26)
18	$4s^24p^4(^1D)4d$	$^2G_{7/2}$	0.37(6)+0.40(8)−0.77(18)+0.29(27)
19	$4s^24p^4(^3P)4d$	$^4D_{7/2}$	0.62(6)+0.65(19)−0.33(8)−0.27(27)
20	$4s^24p^4(^1D)4d$	$^2P_{3/2}$	−0.61(12)−0.74(20)
21	$4s^24p^4(^3P)4d$	$^2D_{5/2}$	0.27(17)+0.28(5)−0.29(11)+0.60(21)−0.43(13)+0.44(26)
22	$4s^24p^4(^3P)4d$	$^2P_{3/2}$	−0.32(60)−0.37(3) 0.69(22)−0.50(16)
23	$4s^24p^4(^1D)4d$	$^2S_{1/2}$	−0.46(4)−0.24(9)−0.49(29)+0.61(23)+0.33(14)
24	$4s^24p^4(^1D)4d$	$^2G_{9/2}$	0.55(10)−0.83(24)
25	$4s^24p^4(^3P)4d$	$^2F_{5/2}$	0.41(5)+0.44(11)+0.67(25)+0.37(26)
26	$4s^24p^4(^1D)4d$	$^2D_{5/2}$	0.49(11)+0.42(21)−0.43(13)−0.59(26)
27	$4s^24p^4(^1D)4d$	$^2F_{7/2}$	−0.34(6)+0.20(19)+0.45(8)−0.25(18)−0.76(27)
28	$4s^24p^4(^3P)4d$	$^2D_{3/2}$	0.36(60)+0.25(3)−0.62(28)−0.52(16)−0.30(20)
29	$4s^24p^4(^1D)4d$	$^2P_{1/2}$	−0.59(9)−0.56(29)−0.53(23)
60	$4s^24p^4(^3P)4d$	$^4F_{3/2}$	−0.45(60)−0.24(3)−0.48(28)−0.36(22)+0.58(7)
66	$4s^24p^4(^1S)4d$	$^2D_{5/2}$	−0.35(17)−0.37(5)−0.36(11)+0.43(25)+0.23(21)+0.60(66)

Table B. Mixing coefficients (MC) for some levels of W XL. Numbers outside and inside a bracket correspond to MC and the level, respectively. See Table 2 for level indices.

Index	Configuration	Level	Mixing coefficients
30	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{3/2}$	0.39(30)+0.25(80)−0.29(101)−0.25(154)−0.21(128)−0.38(54) +0.35(217)−0.24(125)+0.22(46)
31	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{1/2}$	−0.48(48)−0.22(71)−0.24(113)−0.30(218)−0.58(31)−0.30(272)
32	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{5/2}$	−0.39(32)−0.26(117)−0.22(163)−0.34(90)−0.24(153)−0.27(95) +0.42(115)−0.20(39)+0.26(230)+0.24(219)
33	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4H^o_{7/2}$	0.30(96)+0.23(227)+0.26(122)−0.35(33)−0.24(250)+0.36(112) −0.24(37)+0.33(43)+0.34(57)
34	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{3/2}$	0.38(34)+0.24(156)−0.23(167)−0.23(70)+0.26(116)−0.35(170) +0.25(237)+0.21(257)+0.32(290)
35	$4s4p^5(^3P)4d$	$^4P^o_{1/2}$	0.53(35)+0.30(52)−0.39(162)−0.37(119)+0.34(245)
36	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4G^o_{9/2}$	−0.47(97)−0.21(127)+0.33(235)+0.22(165)−0.53(36)+0.30(55)−0.20(254)
37	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{7/2}$	−0.30(96)−0.30(91)−0.22(112)+0.33(37)+0.26(284)−0.23(264)+0.24(57)
38	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4P^o_{3/2}$	−0.29(89)+0.37(58)−0.25(30)+0.23(114)−0.22(108)−0.22(38) −0.38(217)+0.23(293)−0.23(170)−0.22(257)
39	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{5/2}$	−0.20(49)+0.28(59)+0.20(32)−0.26(117)−0.25(90)−0.24(45) +0.35(39)+0.33(230)+0.20(330)
40	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{5/2}$	−0.50(98)+0.23(251)+0.30(132)−0.41(40)+0.44(238)
41	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{7/2}$	−0.24(79)+0.32(41)+0.26(84)−0.24(149)+0.39(264)−0.26(155)+0.27(186)
42	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{11/2}$	0.40(42)+0.37(136)−0.21(103)+0.24(145)−0.33(56)+0.49(239)+0.47(255)
43	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2G^o_{7/2}$	−0.21(68)−0.28(96)−0.35(33)+0.30(37)−0.22(147)+0.36(43)
44	$4s^2 4p^4(^1D)4f$	$^2H^o_{9/2}$	0.26(176)+0.31(73)+0.28(44)−0.39(270)−0.35(179)−0.22(142) −0.29(121)+0.44(157)
45	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4F^o_{5/2}$	−0.20(59)+0.27(32)−0.20(274)−0.23(159)−0.28(45)+0.27(251) −0.23(115)+0.26(230)−0.22(40)−0.26(263)
46	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2P^o_{3/2}$	0.20(30)−0.25(100)−0.29(80)−0.20(101)−0.21(146)+0.22(128) +0.22(126)−0.28(54)+0.21(237)+0.30(318)+0.25(125)−0.30(46)
47	$4s^2 4p^4(^1D)4f$	$^2D^o_{3/2}$	0.24(171)−0.28(150)+0.26(47)+0.28(76)−0.30(89)+0.27(87) −0.25(114)+0.23(217)−0.24(293)
48	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{1/2}$	0.28(61)+0.26(88)−0.59(35)−0.25(48)+0.21(71) −0.23(31)+0.27(272)
49	$4s4p^5(^3P)4d$	$^4F^o_{5/2}$	−0.35(151)+0.22(72)+0.28(187)−0.42(49)+0.24(59) +0.25(0)−0.26(85)−0.21(152)−0.33(330)
50	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2F^o_{5/2}$	−0.21(168)−0.23(53)−0.23(67)−0.27(163)−0.29(50) −0.21(324)+0.35(219)
51	$4s4p^5(^3P)4d$	$^4F^o_{7/2}$	0.38(68)+0.56(51)−0.25(75)+0.31(79)
52	$4s^2 4p^3(^4S)4d^2(^3P)$	$^4P^o_{1/2}$	0.45(35)+0.32(189)−0.35(52)−0.21(106)−0.30(62) +0.24(119)+0.28(306)
53	$4s^2 4p^4(^1S)4f$	$^2F^o_{5/2}$	−0.20(151)−0.28(53)+0.23(77)−0.22(159)+0.31(85) +0.26(152)−0.21(238)−0.21(263)+0.39(330)
54	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{3/2}$	−0.22(150)−0.34(89)+0.33(58)−0.35(34)−0.30(167) −0.29(237)+0.32(257)
55	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{9/2}$	0.40(65)+0.24(184)+0.63(74)−0.27(55)−0.22(254)
56	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2I^o_{11/2}$	0.39(69)+0.27(175)+0.37(42)−0.31(136)+0.20(145) +0.30(56)+0.43(239)−0.42(255)
57	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2F^o_{7/2}$	0.24(325)+0.26(75)+0.28(79)−0.21(227)+0.27(41) +0.22(122)−0.23(250)+0.23(147)+0.31(264)+0.32(57)
58	$4s4p^5(^3P)4d$	$^4P^o_{3/2}$	0.25(171)−0.25(63)+0.29(47)−0.56(58)+0.33(87)
59	$4s4p^5(^3P)4d$	$^4D^o_{5/2}$	0.24(168)+0.27(53)+0.48(59)−0.26(67)+0.25(77)−0.37(78)
61	$4s^2 4p^4(^3P)4f$	$^4D^o_{1/2}$	0.54(61)+0.40(148)+0.34(88)+0.26(189)−0.28(272)
62	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4D^o_{1/2}$	−0.22(86)−0.29(130)−0.28(71)−0.41(62)−0.32(162) +0.40(306)+0.28(245)−0.32(272)
63	$4s^2 4p^4(^3P)4f$	$^2D^o_{3/2}$	0.36(150)−0.44(63)+0.43(161)-0.21(83)+0.34(188)
64	$4s^2 4p^4(^3P)4f$	$^2F^o_{7/2}$	−0.27(68)+0.22(185)+0.38(64)+0.28(164)+0.26(234) −0.21(250)−0.21(147)−0.25(284)+0.24(57)
65	$4s^2 4p^4(^3P)4f$	$^4F^o_{9/2}$	−0.27(176)+0.41(65)+0.29(184)−0.27(97)+0.24(127) −0.27(235)−0.29(92)+0.37(55)+0.34(254)

Table C. Comparison of excitation energies (in Ryd) for some levels of W XL. Level orderings correspond to those of Table 2.

Index	Configuration	Level	GRASP	FAC1	FAC2
30	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{3/2}$	22.5740	22.5805	22.5916
31	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{1/2}$	22.7106	22.7166	22.7278
32	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{5/2}$	22.7488	22.7545	22.7658
33	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4H^o_{7/2}$	22.8911	22.8951	22.9063
34	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{3/2}$	23.3766	23.3797	23.3891
35	$4s4p^5(^3P)4d$	$^4P^o_{1/2}$	24.0349	24.0451	24.0555
36	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4G^o_{9/2}$	24.0548	24.0666	24.0781
37	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{7/2}$	24.0662	24.0779	24.0890
38	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4P^o_{3/2}$	24.0928	24.1036	24.1143
39	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{5/2}$	24.2275	24.2371	24.2480
40	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{5/2}$	24.3977	24.4083	24.4189
41	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{7/2}$	24.4336	24.4426	24.4532
42	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{11/2}$	24.5388	24.5484	24.5595
43	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2G^o_{7/2}$	24.9901	24.9957	25.0069
44	$4s^2 4p^4(^1D)4f$	$^2H^o_{9/2}$	24.9948	25.0028	25.0136
45	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4F^o_{5/2}$	25.0279	25.0330	25.0439
46	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2P^o_{3/2}$	25.2438	25.2483	25.2591
47	$4s^2 4p^4(^1D)4f$	$^2D^o_{3/2}$	25.2599	25.2588	25.2689
48	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{1/2}$	25.2688	25.2699	25.2803
49	$4s4p^5(^3P)4d$	$^4F^o_{5/2}$	25.2857	25.2857	25.2958
50	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2F^o_{5/2}$	25.4722	25.4819	25.4926
51	$4s4p^5(^3P)4d$	$^4F^o_{7/2}$	25.4883	25.4873	25.4972
52	$4s^2 4p^3(^4S)4d^2(^3P)$	$^4P^o_{1/2}$	25.5114	25.5113	25.5215
53	$4s^2 4p^4(^1S)4f$	$^2F^o_{5/2}$	25.6214	25.6295	25.6399
54	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{3/2}$	25.7739	25.7706	25.7807
55	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{9/2}$	25.8007	25.8092	25.8194
56	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2I^o_{11/2}$	25.8381	25.8518	25.8626
57	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2F^o_{7/2}$	25.9381	25.9510	25.9616
58	$4s4p^5(^3P)4d$	$^4P^o_{3/2}$	26.0483	26.0447	26.0545
59	$4s4p^5(^3P)4d$	$^4D^o_{5/2}$	26.3546	26.3458	26.3550
61	$4s^2 4p^4(^3P)4f$	$^4D^o_{1/2}$	26.4363	26.4309	26.4398
62	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4D^o_{1/2}$	26.7300	26.7355	26.7458
63	$4s^2 4p^4(^3P)4f$	$^2D^o_{3/2}$	26.7611	26.7650	26.7740
64	$4s^2 4p^4(^3P)4f$	$^2F^o_{7/2}$	26.7815	26.7908	26.8009
65	$4s^2 4p^4(^3P)4f$	$^4F^o_{9/2}$	26.8418	26.8461	26.8566

GRASP: present calculations from the GRASP code with 4215 levels
FAC1: present calculations from the FAC code with 4215 levels
FAC2: present calculations from the FAC code with 11,849 levels

Table D. Comparison of oscillator strengths (f- values) for some transitions of W XL. See Table 2 for level indices. $a\pm b \equiv a\times 10^{\pm b}$.

I	J	GRASP	FAC1	FAC2	Ratio	I	J	GRASP	FAC1	FAC2	Ratio
3	30	3.9112-3	3.926-3	3.927-3	1.1-0	4	52	7.6985-3	5.016-3	5.018-3	7.5-1
3	31	1.4208-3	1.396-3	1.396-3	1.1-0	4	54	1.0811-1	1.066-1	1.068-1	1.1-0
3	34	1.1878-5	1.145-5	1.135-5	1.5-0	4	58	4.5795-2	4.783-2	4.791-2	1.1-0
3	35	9.0205-4	5.882-4	5.849-4	1.9-0	4	61	1.4751-2	1.145-2	1.155-2	1.9-0
3	38	1.0258-2	9.486-3	9.458-3	1.3-0	4	62	1.8444-2	1.725-2	1.722-2	1.0-0
3	39	2.7366-2	2.587-2	2.575-2	1.2-0	4	63	3.4014-3	2.880-3	2.923-3	9.2-1
3	40	3.7872-4	3.306-4	3.293-4	9.1-1	5	30	2.0459-3	2.051-3	2.054-3	1.5-0
3	45	1.9742-1	2.048-1	2.048-1	8.3-1	5	32	1.0676-4	1.194-4	1.204-4	6.6-1
3	46	7.8150-2	7.612-2	7.557-2	9.8-1	5	33	2.5187-3	2.633-3	2.630-3	1.4-0
3	47	5.9716-3	9.911-3	1.048-2	6.7-1	5	34	4.0929-3	3.965-3	3.961-3	1.3-0
3	48	2.7463-2	3.123-2	3.126-2	7.4-1	5	37	8.3477-5	6.747-5	6.851-5	4.3-0
3	49	2.6154-3	2.169-3	2.066-3	8.9-2	5	38	1.2041-3	9.124-4	9.082-4	9.8-1
3	50	2.5621-2	2.616-2	2.624-2	9.6-1	5	39	2.6589-3	2.290-3	2.285-3	1.6-0
3	52	3.9984-2	3.936-2	3.937-2	8.3-1	5	40	6.4753-5	5.028-5	5.096-5	2.9-0
3	53	1.2948-2	1.173-2	1.174-2	1.2-0	5	41	1.2505-3	1.130-3	1.122-3	1.7-0
3	54	6.0372-2	6.791-2	6.780-2	6.8-1	5	43	1.9014-1	1.950-1	1.947-1	8.5-1
3	58	4.9802-2	4.512-2	4.522-2	1.2-0	5	45	4.4200-2	4.139-2	4.131-2	1.0-0
3	59	2.5836-4	4.468-4	4.464-4	9.4-1	5	46	8.9965-3	9.916-3	9.895-3	1.0-0
3	61	4.6960-3	5.704-3	5.667-3	3.1-1	5	47	1.3296-4	9.793-5	1.170-4	1.9-0
3	62	3.6121-5	1.570-5	1.588-5	5.8-0	5	49	1.8391-2	2.491-2	2.509-2	6.0-1
3	63	4.0475-3	3.542-3	3.575-3	9.3-1	5	50	2.2852-2	2.426-2	2.421-2	6.9-1
4	30	8.3655-3	8.354-3	8.344-3	1.3-0	5	51	1.6155-2	1.480-2	1.495-2	1.5-0
4	31	1.8952-2	1.884-2	1.882-2	1.2-0	5	53	1.5529-2	1.468-2	1.458-2	5.6-1
4	34	2.1947-4	1.880-4	1.847-4	3.2-2	5	54	6.2702-2	6.808-2	6.809-2	9.3-1
4	35	3.2561-3	2.644-3	2.639-3	1.7-0	5	57	4.1634-4	3.568-4	3.562-4	6.0-1
4	38	4.4388-3	3.927-3	3.907-3	1.7-0	5	58	4.8795-2	4.782-2	4.779-2	6.4-1
4	46	1.9806-1	2.110-1	2.143-1	8.3-1	5	59	6.6341-2	6.831-2	6.829-2	9.1-1
4	47	1.2180-1	1.158-1	1.124-1	7.6-1	5	63	3.2631-3	3.325-3	3.354-3	8.2-1
4	48	6.0750-2	6.826-2	6.827-2	8.3-1	5	64	9.6953-3	9.421-3	9.327-3	9.1-1

GRASP: present calculations from the GRASP code with 4215 levels
FAC1: present calculations from the FAC code with 4215 levels
FAC2: present calculations from the FAC code with 11,849 levels
Ratio: ratio of velocity and length forms of f- values from the GRASP code

Explanation of Tables

Table 1. Configurations and levels of W XL.

Index	Configuration Index
Configuration	The configuration included in the calculation
No. of Levels	The number of levels the configuration generates. Odd parity levels are designated with a superscript "o"
Energy Range	Energy range of the levels in Ryd unit
GRASPa	earlier calculations [5] from the GRASP code with 4128 levels
GRASPb	present calculations from the GRASP code with 4215 levels
RELAC	Fournier [1]
Y	If a configuration is included under a calculation

Table 2. Energies (Ryd) for the lowest 360 levels of W XL and their lifetimes (τ, s).

Index	Level Index
Configuration	The configuration to which the level belongs
Level	The LSJ designation of the level
Energy	Present energies from the GRASP code with 4215 level calculations
τ (s)	Lifetime of the level in s

Table 3. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units) for electric dipole (E1), and A_{ji} for electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions of W XL.

i and j	The lower (i) and upper (j) levels of a transition as defined in Table 2.
λ_{ij}	Transition wavelength (in Å)
A_{ji}^{E1}	Radiative transition probability (in s^{-1}) for the E1 transitions
f_{ij}^{E1}	Absorption oscillator strength (dimensionless) for the E1 transitions
S^{E1}	Line strength in atomic unit (a.u.), 1 a.u. = 6.460×10^{-36} cm^2 esu^2 for the E1 transitions
A_{ji}^{E2}	Radiative transition probability (in s^{-1}) for the E2 transitions
A_{ji}^{M1}	Radiative transition probability (in s^{-1}) for the M1 transitions
A_{ji}^{M2}	Radiative transition probability (in s^{-1}) for the M2 transitions
$a\pm b$	$\equiv a \times 10^{\pm b}$

Table 1. Configurations and levels of W XL.

Index	Configuration	No. of Levels	Energy Range (Ryd)	GRASPa	GRASPb	RELAC
1	$4s^2 4p^5$	2^o	0–7	Y	Y	Y
2	$4s^2 4p^4 4d$	28	11–27	Y	Y	Y
3	$4s^2 4p^4 4f$	30^o	25–42	Y	Y	Y
4	$4s4p^6$	1	15	Y	Y	Y
5	$4p^6 4d$	2	43–45	Y	Y	
6	$4p^6 4f$	2^o	58–59	Y	Y	
7	$4s4p^5 4d$	23^o	25–36	Y	Y	Y
8	$4s4p^5 4f$	24	41–50	Y	Y	Y
9	$4s^2 4p^3 4d^2$	141^o	22–42	Y	Y	
10	$4s^2 4p^3 4f^2$	221^o	52–71	Y	Y	
11	$4s^2 4p^3 4d4f$	363	37–58	Y	Y	
12	$4s^2 4p^2 4d^3$	261	34–57	Y	Y	
13	$4s^2 4p4d^4$	180^o	54–73	Y	Y	
14	$4s^2 4p^2 4d^2 4f$	1140^o	49–73	Y	Y	
15	$4s4p^3 4d^3$	678^o	47–73	Y	Y	
16	$4p^5 4d^2$	45^o	53–64	Y	Y	
17	$3d^9 4s^2 4p^5 4d$	96^o	131–147	Y	Y	Y
18	$3d^9 4s^2 4p^5 4f$	113	147–161	Y	Y	Y
19	$3d^9 4s^2 4p^6$	2	120–126	Y	Y	Y
20	$3p^5 3d^{10} 4s^2 4p^6$	2^o	155–178	Y	Y	Y
21	$3p^5 3d^{10} 4s^2 4p^5 4d$	65	165–198	Y	Y	Y
22	$3p^5 3d^{10} 4s^2 4p^5 4f$	36^o	180–212	Y	Y	Y
23	$4s4p^5 5s$	7^o	65–74	Y	Y	
24	$4s4p^5 5p$	18	68–80	Y	Y	
25	$4s4p^5 5d$	23^o	77–86	Y	Y	
26	$4s4p^5 5f$	24	84–92	Y	Y	
27	$4s4p^5 5g$	24^o	87–96	Y	Y	
28	$4p^6 5s$	1	82	Y	Y	
29	$4p^6 5p$	2^o	86–89	Y	Y	
30	$4p^6 5d$	2	94–95	Y	Y	
31	$4p^6 5f$	2^o	100–101	Y	Y	
32	$4p^6 5g$	2	104–105	Y	Y	
33	$4s^2 4p^4 5s$	8	50–65	Y	Y	Y
34	$4s^2 4p^4 5p$	21^o	54–72	Y	Y	Y
35	$4s^2 4p^4 5d$	28	62–78	Y	Y	Y
36	$4s^2 4p^4 5f$	30^o	69–84	Y	Y	Y
37	$4s^2 4p^4 5g$	30	73–88	Y	Y	
38	$3d^9 4s^2 4p^5 5s$	23^o	172–185	Y	Y	Y
39	$3d^9 4s^2 4p^5 5p$	65	175–191	Y	Y	Y
40	$3d^9 4s^2 4p^5 5d$	96^o	183–197	Y	Y	Y
41	$3d^9 4s^2 4p^5 5f$	113	190–203	Y	Y	Y
42	$3d^9 4s^2 4p^5 5g$	119^o	194–207	Y	Y	
43	$4s^2 4p^4 6s$	8	81–96		Y	
44	$4s^2 4p^4 6p$	21^o	81–99		Y	
45	$4s^2 4p^4 6d$	28	87–102		Y	
46	$4s^2 4p^4 6f$	30^o	91–106		Y	

GRASPa: earlier calculations [5] from the GRASP code with 4128 levels
GRASPb: present calculations from the GRASP code with 4215 levels
RELAC: Fournier [1]
Y: configuration included under a calculation

Table 2. Energies (Ryd) for the lowest 360 levels of W XL and their lifetimes (τ, s). $a\pm b \equiv a\times 10^{\pm b}$.

Index	Configuration	Level	Energy	τ (s)
1	$4s^2 4p^5$	$^2P^o_{3/2}$	0.0000	.....
2	$4s^2 4p^5$	$^2P^o_{1/2}$	6.7987	1.314-07
3	$4s^2 4p^4(^3P)4d$	$^4D_{3/2}$	11.2603	3.216-10
4	$4s^2 4p^4(^3P)4d$	$^4P_{1/2}$	11.4314	7.153-11
5	$4s^2 4p^4(^3P)4d$	$^4D_{5/2}$	11.4275	6.342-10
6	$4s^2 4p^4(^3P)4d$	$^4F_{7/2}$	11.6140	4.757-03
7	$4s^2 4p^4(^1S)4d$	$^2D_{3/2}$	12.1537	2.679-09
8	$4s^2 4p^4(^3P)4d$	$^2F_{7/2}$	12.7500	2.237-05
9	$4s^2 4p^4(^3P)4d$	$^2P_{1/2}$	12.8178	1.040-10
10	$4s^2 4p^4(^3P)4d$	$^4F_{9/2}$	12.8147	3.704-05
11	$4s^2 4p^4(^3P)4d$	$^4P_{5/2}$	13.7448	3.062-10
12	$4s^2 4p^4(^3P)4d$	$^4P_{3/2}$	14.0956	2.361-12
13	$4s^2 4p^4(^1D)4d$	$^2F_{5/2}$	14.3278	1.752-12
14	$4s4p^6$	$^2S_{1/2}$	15.3699	1.656-12
15	$4s^2 4p^4(^3P)4d$	$^4D_{1/2}$	17.6737	1.220-10
16	$4s^2 4p^4(^1D)4d$	$^2D_{3/2}$	18.0340	5.114-10
17	$4s^2 4p^4(^3P)4d$	$^4F_{5/2}$	18.3401	2.068-11
18	$4s^2 4p^4(^1D)4d$	$^2G_{7/2}$	18.4099	2.604-07
19	$4s^2 4p^4(^3P)4d$	$^4D_{7/2}$	19.3377	1.356-07
20	$4s^2 4p^4(^1D)4d$	$^2P_{3/2}$	19.6012	1.315-12
21	$4s^2 4p^4(^3P)4d$	$^2D_{5/2}$	19.7541	5.212-13
22	$4s^2 4p^4(^3P)4d$	$^2P_{3/2}$	19.7592	1.001-12
23	$4s^2 4p^4(^1D)4d$	$^2S_{1/2}$	19.8287	3.712-13
24	$4s^2 4p^4(^1D)4d$	$^2G_{9/2}$	19.7982	2.445-07
25	$4s^2 4p^4(^3P)4d$	$^2F_{5/2}$	19.8152	1.731-11
26	$4s^2 4p^4(^1D)4d$	$^2D_{5/2}$	20.0716	1.664-11
27	$4s^2 4p^4(^1D)4d$	$^2F_{7/2}$	20.3657	2.046-07
28	$4s^2 4p^4(^3P)4d$	$^2D_{3/2}$	21.3920	1.224-12
29	$4s^2 4p^4(^1D)4d$	$^2P_{1/2}$	21.8225	7.443-13
30	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{3/2}$	22.5740	7.837-11
31	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{1/2}$	22.7106	3.897-11
32	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{5/2}$	22.7488	8.513-11
33	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4H^o_{7/2}$	22.8911	2.090-10
34	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{3/2}$	23.3766	1.134-10
35	$4s4p^5(^3P)4d$	$^4P^o_{1/2}$	24.0349	6.245-11
36	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4G^o_{9/2}$	24.0548	1.186-10
37	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{7/2}$	24.0662	6.165-11
38	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4P^o_{3/2}$	24.0928	3.518-11
39	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{5/2}$	24.2275	2.461-11
40	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{5/2}$	24.3977	1.774-10
41	$4s^2 4p^3(^4S)4d^2(^3P)$	$^6P^o_{7/2}$	24.4336	5.739-11
42	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{11/2}$	24.5388	3.953-08
43	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2G^o_{7/2}$	24.9901	3.984-12
44	$4s^2 4p^4(^1D)4f$	$^2H^o_{9/2}$	24.9948	7.327-12
45	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4F^o_{5/2}$	25.0279	3.599-12
46	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2P^o_{3/2}$	25.2438	3.099-12
47	$4s^2 4p^4(^1D)4f$	$^2D^o_{3/2}$	25.2599	7.771-12
48	$4s^2 4p^3(^4S)4d^2(^3F)$	$^6F^o_{1/2}$	25.2688	4.826-12
49	$4s4p^5(^3P)4d$	$^4F^o_{5/2}$	25.2857	1.259-11
50	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2F^o_{5/2}$	25.4722	8.736-12

Index	Configuration	Level	Energy	τ (s)
51	$4s4p^5(^3P)4d$	$^4F^o_{7/2}$	25.4883	6.138-12
52	$4s^24p^3(^4S)4d^2(^3P)$	$^4P^o_{1/2}$	25.5114	4.364-12
53	$4s^24p^4(^1S)4f$	$^2F^o_{5/2}$	25.6214	9.985-12
54	$4s^24p^3(^2P)4d^2(^3F)$	$^4F^o_{3/2}$	25.7739	2.222-12
55	$4s^24p^3(^2P)4d^2(^3F)$	$^4F^o_{9/2}$	25.8007	2.075-11
56	$4s^24p^3(^2D)4d^2(^1G)$	$^2I^o_{11/2}$	25.8381	3.685-11
57	$4s^24p^3(^2P)4d^2(^1D)$	$^2F^o_{7/2}$	25.9381	2.762-11
58	$4s4p^5(^3P)4d$	$^4P^o_{3/2}$	26.0483	2.805-12
59	$4s4p^5(^3P)4d$	$^4D^o_{5/2}$	26.3546	2.217-12
60	$4s^24p^4(^3P)4d$	$^4F_{3/2}$	26.2847	6.222-13
61	$4s^24p^4(^3P)4f$	$^4D^o_{1/2}$	26.4363	2.843-12
62	$4s^24p^3(^2D)4d^2(^3P)$	$^4D^o_{1/2}$	26.7300	1.943-12
63	$4s^24p^4(^3P)4f$	$^2D^o_{3/2}$	26.7611	9.961-12
64	$4s^24p^4(^3P)4f$	$^2F^o_{7/2}$	26.7815	3.258-12
65	$4s^24p^4(^3P)4f$	$^4F^o_{9/2}$	26.8418	1.687-12
66	$4s^24p^4(^1S)4d$	$^2D_{5/2}$	26.8956	4.372-10
67	$4s4p^5(^3P)4d$	$^4P^o_{5/2}$	27.1442	3.084-12
68	$4s^24p^4(^3P)4f$	$^4F^o_{7/2}$	27.2070	1.151-12
69	$4s^24p^4(^3P)4f$	$^4G^o_{11/2}$	27.2070	2.458-12
70	$4s^24p^3(^2D)4d^2(^1S)$	$^2D^o_{3/2}$	27.2154	3.049-12
71	$4s^24p^3(^4S)4d^2(^1D)$	$^4D^o_{1/2}$	27.2956	1.246-12
72	$4s^24p^4(^3P)4f$	$^4D^o_{5/2}$	27.3080	1.209-12
73	$4s^24p^4(^3P)4f$	$^2G^o_{9/2}$	27.3660	1.250-12
74	$4s4p^5(^3P)4d$	$^4F^o_{9/2}$	27.4088	2.004-12
75	$4s4p^5(^3P)4d$	$^4D^o_{7/2}$	27.4358	2.953-12
76	$4s4p^5(^3P)4d$	$^4F^o_{3/2}$	27.4611	1.206-12
77	$4s4p^5(^3P)4d$	$^2D^o_{5/2}$	27.6888	3.862-12
78	$4s4p^5(^1P)4d$	$^2F^o_{5/2}$	27.8081	1.012-12
79	$4s4p^5(^3P)4d$	$^2F^o_{7/2}$	27.8172	3.989-12
80	$4s^24p^3(^4S)4d^2(^1D)$	$^4D^o_{3/2}$	27.8684	1.664-12
81	$4s4p^5(^1P)4d$	$^2F^o_{7/2}$	28.2852	1.876-12
82	$4s^24p^4(^3P)4f$	$^2D^o_{5/2}$	28.3289	1.442-12
83	$4s4p^5(^3P)4d$	$^2D^o_{3/2}$	28.3361	1.288-12
84	$4s^24p^3(^4S)4d^2(^1G)$	$^4G^o_{7/2}$	28.7019	9.732-13
85	$4s^24p^3(^4S)4d^2(^1G)$	$^4G^o_{5/2}$	28.7824	1.350-12
86	$4s4p^5(^3P)4d$	$^2P^o_{1/2}$	29.1819	8.802-13
87	$4s4p^5(^1P)4d$	$^2D^o_{3/2}$	29.3774	8.999-13
88	$4s4p^5(^3P)4d$	$^4D^o_{1/2}$	29.5049	6.931-13
89	$4s4p^5(^3P)4d$	$^4D^o_{3/2}$	29.5002	8.104-13
90	$4s^24p^3(^2D)4d^2(^3F)$	$^4G^o_{5/2}$	29.5211	9.909-13
91	$4s^24p^3(^2D)4d^2(^3F)$	$^4G^o_{7/2}$	29.7172	1.044-12
92	$4s^24p^3(^2D)4d^2(^3F)$	$^4G^o_{9/2}$	29.9721	1.326-12
93	$4s^24p^3(^2D)4d^2(^3P)$	$^4P^o_{3/2}$	30.2778	4.493-12
94	$4s^24p^3(^2D)4d^2(^3P)$	$^4P^o_{5/2}$	30.2981	4.743-12
95	$4s^24p^3(^2P)4d^2(^3F)$	$^4G^o_{5/2}$	30.3400	1.148-12
96	$4s^24p^3(^4S)4d^2(^3F)$	$^6F^o_{7/2}$	30.3322	7.388-12
97	$4s^24p^3(^4S)4d^2(^3F)$	$^6F^o_{9/2}$	30.3904	1.623-11
98	$4s^24p^3(^4S)4d^2(^3P)$	$^6P^o_{5/2}$	30.4657	2.926-11
99	$4s^24p^3(^2D)4d^2(^3P)$	$^4F^o_{7/2}$	30.5708	3.148-11
100	$4s^24p^3(^4S)4d^2(^3P)$	$^4P^o_{3/2}$	30.6237	7.656-13

Index	Configuration	Level	Energy	τ (s)
101	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4F^o_{3/2}$	30.6638	3.488-12
102	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2F^o_{5/2}$	30.6715	6.051-12
103	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4H^o_{11/2}$	30.7200	1.403-10
104	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4F^o_{9/2}$	30.7486	1.461-11
105	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4D^o_{5/2}$	30.9644	7.656-13
106	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4P^o_{1/2}$	31.0257	3.548-13
107	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2H^o_{11/2}$	31.0384	1.748-10
108	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4D^o_{3/2}$	31.1739	3.740-13
109	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4D^o_{5/2}$	31.2920	3.609-13
110	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2F^o_{7/2}$	31.3150	3.785-13
111	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2I^o_{13/2}$	31.2886	7.837-07
112	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4G^o_{7/2}$	31.4697	9.805-12
113	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4D^o_{1/2}$	31.5175	7.772-13
114	$4s^2 4p^3(^4S)4d^2(^3F)$	$^4F^o_{3/2}$	31.5670	2.035-12
115	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4F^o_{5/2}$	31.5925	3.505-12
116	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{3/2}$	31.6625	5.767-12
117	$4s^2 4p^3(^4S)4d^2(^3F)$	$^4F^o_{5/2}$	31.6849	4.289-12
118	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4G^o_{11/2}$	31.6619	4.508-12
119	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4P^o_{1/2}$	31.7827	9.973-12
120	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2G^o_{7/2}$	31.8142	2.127-12
121	$4s^2 4p^3(^2P)4d^2(^1G)$	$^2H^o_{9/2}$	31.8419	1.851-12
122	$4s^2 4p^3(^4S)4d^2(^1D)$	$^4D^o_{7/2}$	31.8587	4.434-12
123	$4s^2 4p^3(^2D)4d^2(^3F)$	$^2H^o_{9/2}$	31.9066	2.598-12
124	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{1/2}$	31.9686	8.495-13
125	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2D^o_{3/2}$	31.9999	3.438-13
126	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2P^o_{3/2}$	32.0297	9.743-12
127	$4s^2 4p^3(^4S)4d^2(^3F)$	$^4F^o_{9/2}$	32.1176	6.846-13
128	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2D^o_{3/2}$	32.1400	1.862-12
129	$4s^2 4p^3(^2D)4d^2(^3F)$	$^2G^o_{7/2}$	32.1537	8.324-13
130	$4s^2 4p^3(^4S)4d^2(^3P)$	$^2P^o_{1/2}$	32.2018	5.667-13
131	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4F^o_{9/2}$	32.1955	1.344-12
132	$4s^2 4p^3(^2D)4d^2(^3P)$	$^2D^o_{5/2}$	32.2157	1.423-12
133	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2D^o_{3/2}$	32.2322	5.690-13
134	$4s^2 4p^3(^2D)4d^2(^3P)$	$^2F^o_{7/2}$	32.2737	8.840-13
135	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4P^o_{5/2}$	32.2854	6.761-13
136	$4s^2 4p^3(^4S)4d^2(^1G)$	$^4G^o_{11/2}$	32.3397	6.554-13
137	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2D^o_{5/2}$	32.3545	1.091-11
138	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2G^o_{9/2}$	32.3781	1.432-12
139	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4D^o_{7/2}$	32.4195	5.067-13
140	$4s^2 4p^3(^2D)4d^2(^3F)$	$^4H^o_{13/2}$	32.4627	7.531-07
141	$4s^2 4p^3(^2D)4d^2(^3P)$	$^2F^o_{5/2}$	32.5074	1.414-12
142	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2G^o_{9/2}$	32.5296	1.873-12
143	$4s^2 4p^3(^2D)4d^2(^1D)$	$^2S^o_{1/2}$	32.6525	7.939-13
144	$4s^2 4p^3(^2D)4d^2(^3F)$	$^2F^o_{7/2}$	32.6311	7.505-12
145	$4s^2 4p^3(^2D)4d^2(^3F)$	$^2H^o_{11/2}$	32.8560	5.634-12
146	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4D^o_{3/2}$	32.9126	8.632-13
147	$4s^2 4p^3(^2P)4d^2(^3F)$	$^4D^o_{7/2}$	32.9583	1.110-12
148	$4s^2 4p^4(^1D)4f$	$^2P^o_{1/2}$	33.0138	9.666-13
149	$4s^2 4p^3(^2D)4d^2(^1G)$	$^2F^o_{7/2}$	33.0228	8.045-13
150	$4s^2 4p^4(^3P)4f$	$^4D^o_{3/2}$	33.0475	9.569-13

Index	Configuration	Level	Energy	τ (s)
151	$4s^24p^4(^3P)4f$	$^4F^o_{5/2}$	33.0857	8.277-13
152	$4s^24p^3(^2D)4d^2(^1G)$	$^2D^o_{5/2}$	33.1122	7.518-13
153	$4s^24p^3(^2D)4d^2(^3F)$	$^2F^o_{5/2}$	33.2444	4.622-13
154	$4s^24p^3(^2D)4d^2(^3F)$	$^2D^o_{3/2}$	33.2493	6.379-13
155	$4s^24p^3(^2P)4d^2(^1G)$	$^2G^o_{7/2}$	33.3410	9.530-13
156	$4s^24p^3(^4S)4d^2(^1S)$	$^4S^o_{3/2}$	33.4244	4.329-12
157	$4s^24p^3(^2P)4d^2(^1G)$	$^2G^o_{9/2}$	33.5108	3.935-11
158	$4s^24p^3(^2D)4d^2(^1D)$	$^2P^o_{1/2}$	33.5491	4.475-13
159	$4s^24p^3(^4S)4d^2(^3P)$	$^4P^o_{5/2}$	33.6177	5.595-13
160	$4s^24p^3(^2P)4d^2(^1D)$	$^2F^o_{5/2}$	33.6898	4.204-12
161	$4s^24p^4(^1D)4f$	$^2P^o_{3/2}$	33.7429	5.540-13
162	$4s^24p^3(^2D)4d^2(^3P)$	$^4P^o_{1/2}$	33.7999	1.418-12
163	$4s^24p^3(^4S)4d^2(^1D)$	$^4D^o_{5/2}$	33.8166	1.938-12
164	$4s^24p^4(^1D)4f$	$^2F^o_{7/2}$	33.8490	1.137-12
165	$4s^24p^3(^2D)4d^2(^3F)$	$^2G^o_{9/2}$	33.8509	1.534-12
166	$4s^24p^3(^2D)4d^2(^3P)$	$^2P^o_{3/2}$	33.9405	1.079-12
167	$4s^24p^3(^2D)4d^2(^3P)$	$^4F^o_{3/2}$	33.9611	5.367-13
168	$4s^24p^4(^3P)4f$	$^2F^o_{5/2}$	34.0812	1.254-12
169	$4s^24p^4(^3P)4f$	$^4G^o_{7/2}$	34.1138	1.191-12
170	$4s^24p^3(^2P)4d^2(^3P)$	$^4P^o_{3/2}$	34.1551	5.995-13
171	$4s^24p^4(^3P)4f$	$^4F^o_{3/2}$	34.2595	5.235-13
172	$4s^24p^3(^2D)4d^2(^1S)$	$^2D^o_{5/2}$	34.2536	1.095-12
173	$4s^24p^3(^2P)4d^2(^1S)$	$^2P^o_{1/2}$	34.2895	1.954-11
174	$4s^24p^4(^3P)4f$	$^2G^o_{7/2}$	34.3142	1.032-12
175	$4s^24p^4(^1D)4f$	$^2H^o_{11/2}$	34.3375	1.654-12
176	$4s^24p^4(^3P)4f$	$^4G^o_{9/2}$	34.3583	7.295-13
177	$4s^24p^4(^3P)4f$	$^4G^o_{5/2}$	34.4085	6.496-13
178	$4s^24p^3(^2D)4d^2(^3P)$	$^2D^o_{3/2}$	34.4506	7.855-13
179	$4s^24p^3(^2D)4d^2(^1G)$	$^2H^o_{9/2}$	34.5211	7.813-13
180	$4s^24p^4(^1D)4f$	$^2G^o_{7/2}$	34.5183	9.573-13
181	$4s^24p^3(^2D)4d^2(^3F)$	$^2D^o_{5/2}$	34.6437	7.712-13
182	$4s4p^5(^3P)4d$	$^2F^o_{5/2}$	34.6813	1.497-12
183	$4s^24p^2(^3P)4d^3(^4F)$	$^6G_{3/2}$	34.7989	1.102-11
184	$4s^24p^4(^1D)4f$	$^2G^o_{9/2}$	34.7912	1.759-12
185	$4s^24p^4(^3P)4f$	$^4D^o_{7/2}$	34.8652	8.329-13
186	$4s^24p^3(^2P)4d^2(^1G)$	$^2F^o_{7/2}$	35.0611	4.362-13
187	$4s^24p^4(^1D)4f$	$^2F^o_{5/2}$	35.1062	4.703-13
188	$4s4p^5(^1P)4d$	$^2P^o_{3/2}$	35.2229	7.874-13
189	$4s4p^5(^1P)4d$	$^2P^o_{1/2}$	35.4454	3.586-13
190	$4s4p^5(^1P)4d$	$^2D^o_{5/2}$	35.4299	7.846-13
191	$4s4p^5(^3P)4d$	$^2P^o_{3/2}$	35.4765	3.672-13
192	$4s^24p^3(^2D)4d^2(^3F)$	$^4D^o_{7/2}$	35.4617	7.643-13
193	$4s^24p^3(^2D)4d^2(^3P)$	$^2P^o_{1/2}$	35.9327	5.424-13
194	$4s^24p^4(^1D)4f$	$^2D^o_{5/2}$	35.9782	6.625-13
195	$4s^24p^2(^3P)4d^3(^4F)$	$^6G_{5/2}$	36.2982	5.322-12
196	$4s^24p^2(^3P)4d^3(^2H)$	$^4I_{9/2}$	36.4768	7.931-12
197	$4s^24p^2(^3P)4d^3(^2G)$	$^4H_{7/2}$	36.5089	6.733-12
198	$4s^24p^2(^3P)4d^3(^4P)$	$^6D_{3/2}$	36.5322	4.238-12
199	$4s^24p^3(^2D)4d^2(^3F)$	$^2P^o_{3/2}$	36.5867	4.492-13
200	$4s^24p^2(^3P)4d^3(^4P)$	$^6D_{1/2}$	36.6707	4.195-12

Index	Configuration	Level	Energy	τ (s)
201	$4s^24p^2(^3P)4d^3(^2F)$	$^4G_{5/2}$	37.2564	6.044-12
202	$4s^24p^3(^2P)4d[^3F]4f$	$^4I_{9/2}$	37.4411	6.450-11
203	$4s^24p^3(^4S)4d[^5D]4f$	$^6H_{7/2}$	37.5443	1.366-11
204	$4s^24p^3(^2P)4d[^3F]4f$	$^4I_{11/2}$	37.7860	3.817-11
205	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{1/2}$	37.8422	4.798-12
206	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{3/2}$	37.8573	4.084-12
207	$4s^24p^3(^4S)4d[^5D]4f$	$^6H_{5/2}$	37.8636	1.825-11
208	$4s^24p^2(^1S)4d^3(^4F)$	$^4F_{7/2}$	37.8375	6.831-12
209	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{5/2}$	37.9573	3.662-12
210	$4s^24p^2(^1S)4d^3(^4F)$	$^4F_{9/2}$	38.0020	6.820-12
211	$4s^24p^3(^4S)4d[^5D]4f$	$^4D_{7/2}$	38.0310	4.664-12
212	$4s^24p^3(^4S)4d[^5D]4f$	$^6H_{11/2}$	38.0908	1.398-11
213	$4s^24p^2(^1S)4d^3(^4P)$	$^4P_{5/2}$	38.1382	6.628-12
214	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{11/2}$	38.2583	5.915-12
215	$4s^24p^2(^3P)4d^3(^2D)$	$^4F_{3/2}$	38.2945	3.624-12
216	$4s^24p^3(^2P)4d[^3D]4f$	$^4G_{9/2}$	38.3513	4.391-12
217	$4s^24p^3(^2P)4d^2(^3F)$	$^4D^o_{3/2}$	38.3066	2.726-13
218	$4s^24p^3(^2D)4d^2(^3F)$	$^2P^o_{1/2}$	38.3192	2.568-13
219	$4s^24p^3(^2P)4d^2(^1D)$	$^2D^o_{5/2}$	38.3313	3.154-13
220	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{7/2}$	38.4400	2.294-12
221	$4s^24p^3(^4S)4d[^5D]4f$	$^4G_{5/2}$	38.4433	4.688-12
222	$4s^24p^2(^3P)4d^3(^2P)$	$^4D_{1/2}$	38.4503	2.946-12
223	$4s^24p^3(^4S)4d[^5D]4f$	$^4G_{7/2}$	38.4693	5.898-12
224	$4s^24p^3(^2P)4d[^3F]4f$	$^4I_{13/2}$	38.5208	1.780-11
225	$4s^24p^3(^2P)4d[^3D]4f$	$^4F_{3/2}$	38.5572	1.930-12
226	$4s^24p^3(^4S)4d[^5D]4f$	$^6G_{5/2}$	38.6031	1.216-12
227	$4s^24p^3(^4S)4d^2(^3F)$	$^4D^o_{7/2}$	38.5104	3.105-13
228	$4s^24p^3(^2P)4d[^3P]4f$	$^4G_{9/2}$	38.5901	4.341-12
229	$4s^24p^3(^2P)4d[^3F]4f$	$^4G_{7/2}$	38.7020	1.598-12
230	$4s^24p^3(^2P)4d^2(^3F)$	$^2F^o_{5/2}$	38.6603	4.416-13
231	$4s^24p^3(^2P)4d[^3F]4f$	$^4G_{9/2}$	38.7616	1.732-12
232	$4s^24p^3(^4S)4d[^5D]4f$	$^6G_{3/2}$	38.8038	1.589-12
233	$4s^24p^3(^4S)4d[^3D]4f$	$^2D_{5/2}$	38.7864	2.338-12
234	$4s^24p^3(^2D)4d^2(^3F)$	$^4F^o_{7/2}$	38.7299	5.175-13
235	$4s^24p^3(^2D)4d^2(^3F)$	$^4H^o_{9/2}$	38.7267	6.001-13
236	$4s^24p^3(^2P)4d[^3P]4f$	$^4D_{7/2}$	38.8646	3.297-12
237	$4s^24p^3(^2P)4d^2(^3P)$	$^4S^o_{3/2}$	38.8715	4.126-13
238	$4s^24p^3(^2P)4d^2(^3P)$	$^4P^o_{5/2}$	38.8800	5.550-13
239	$4s^24p^3(^2P)4d^2(^3F)$	$^4G^o_{11/2}$	38.8808	8.715-13
240	$4s^24p^3(^4S)4d[^5D]4f$	$^6D_{3/2}$	39.0236	1.688-12
241	$4s^24p^3(^4S)4d^2(^3P)$	$^2P^o_{3/2}$	38.9353	3.453-13
242	$4s^24p^3(^4S)4d[^5D]4f$	$^6P_{5/2}$	39.0436	1.543-12
243	$4s^24p^3(^4S)4d[^5D]4f$	$^6D_{1/2}$	39.1339	1.635-12
244	$4s^24p^3(^4S)4d^2(^3F)$	$^2F^o_{7/2}$	39.0268	6.102-13
245	$4s^24p^3(^2P)4d^2(^3P)$	$^2S^o_{1/2}$	39.0768	6.573-13
246	$4s^24p^3(^2P)4d[^3D]4f$	$^2F_{5/2}$	39.2277	3.401-12
247	$4s^24p^3(^2D)4d[^3G]4f$	$^4K_{11/2}$	39.3029	2.971-12
248	$4s^24p^3(^2P)4d[^3P]4f$	$^4G_{7/2}$	39.3671	2.568-12
249	$4s^24p^3(^4S)4d[^5D]4f$	$^6F_{9/2}$	39.3325	8.911-12
250	$4s^24p^3(^2D)4d^2(^1D)$	$^2G^o_{7/2}$	39.2706	7.968-13

Index	Configuration	Level	Energy	τ (s)
251	$4s^2 4p^3(^2D)4d^2(^3P)$	$^4F^o_{5/2}$	39.3037	1.243-12
252	$4s^2 4p^3(^4S)4d[^5D]4f$	$^4G_{9/2}$	39.4118	1.178-12
253	$4s^2 4p^2(^3P)4d^3(^2D)$	$^4F_{3/2}$	39.4304	2.623-12
254	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2G^o_{9/2}$	39.3570	1.219-12
255	$4s^2 4p^3(^2P)4d^2(^1G)$	$^2H^o_{11/2}$	39.3478	2.147-12
256	$4s^2 4p^3(^2P)4d[^3P]4f$	$^4D_{1/2}$	39.4918	2.068-12
257	$4s^2 4p^3(^2P)4d^2(^3P)$	$^2D^o_{3/2}$	39.4184	6.135-13
258	$4s^2 4p^3(^2P)4d[^3P]4f$	$^2G_{7/2}$	39.5145	1.903-12
259	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4I_{15/2}$	39.4762	2.918-07
260	$4s^2 4p^3(^2P)4d[^3P]4f$	$^4G_{11/2}$	39.5014	4.190-12
261	$4s^2 4p^3(^2P)4d[^3P]4f$	$^4F_{3/2}$	39.5543	1.869-12
262	$4s^2 4p^3(^2P)4d[^3F]4f$	$^2G_{9/2}$	39.5642	1.429-12
263	$4s^2 4p^3(^2P)4d^2(^3P)$	$^2D^o_{5/2}$	39.4699	1.112-12
264	$4s^2 4p^3(^2P)4d^2(^3P)$	$^4D^o_{7/2}$	39.4742	2.381-12
265	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4G_{11/2}$	39.6604	2.247-12
266	$4s^2 4p^3(^2P)4d[^1D]4f$	$^2P_{3/2}$	39.6948	2.635-12
267	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4F_{7/2}$	39.7084	2.070-12
268	$4s^2 4p^3(^4S)4d[^5D]4f$	$^4D_{1/2}$	39.7984	2.276-12
269	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4D_{5/2}$	39.8252	2.897-12
270	$4s^2 4p^3(^4S)4d^2(^1G)$	$^4G^o_{9/2}$	39.7327	1.339-12
271	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4H_{13/2}$	39.8222	2.067-12
272	$4s^2 4p^3(^2P)4d^2(^1D)$	$^2P^o_{1/2}$	39.7796	1.581-11
273	$4s4p^5(^3P)4f$	$^4F_{5/2}$	39.9006	1.668-12
274	$4s^2 4p^3(^4S)4d^2(^3F)$	$^2F^o_{5/2}$	39.8304	8.526-13
275	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4P_{3/2}$	39.9339	2.968-12
276	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4F_{9/2}$	39.9829	1.624-12
277	$4s^2 4p^3(^2P)4d[^3P]4f$	$^2F_{5/2}$	39.9930	2.012-12
278	$4s4p^5(^3P)4f$	$^4F_{7/2}$	40.0810	1.643-12
279	$4s^2 4p^3(^2P)4d[^3P]4f$	$^2D_{5/2}$	40.1167	1.256-12
280	$4s^2 4p^3(^4S)4d[^5D]4f$	$^6P_{3/2}$	40.1715	2.774-12
281	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4D_{7/2}$	40.2315	2.670-12
282	$4s^2 4p^3(^2P)4d[^3F]4f$	$^2H_{11/2}$	40.2448	2.491-12
283	$4s^2 4p^3(^2P)4d[^3F]4f$	$^4P_{5/2}$	40.2846	3.472-12
284	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2F^o_{7/2}$	40.2252	1.626-11
285	$4s^2 4p^3(^2P)4d[^1D]4f$	$^2G_{9/2}$	40.3648	1.816-12
286	$4s^2 4p^3(^2P)4d[^3P]4f$	$^2F_{7/2}$	40.4118	1.823-12
287	$4s^2 4p^3(^4S)4d[^3D]4f$	$^4D_{3/2}$	40.6137	1.391-12
288	$4s^2 4p^3(^2P)4d[^3F]4f$	$^2I_{13/2}$	40.6099	2.563-12
289	$4s^2 4p^3(^2P)4d[^3P]4f$	$^2G_{9/2}$	40.6484	1.848-12
290	$4s^2 4p^3(^2P)4d^2(^1S)$	$^2P^o_{3/2}$	40.5814	3.244-12
291	$4s^2 4p^3(^2P)4d[^1P]4f$	$^2F_{5/2}$	40.7483	1.288-12
292	$4s^2 4p^3(^2P)4d[^3D]4f$	$^2F_{7/2}$	40.7761	1.115-12
293	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2D^o_{3/2}$	40.7061	4.150-13
294	$4s^2 4p^3(^2P)4d[^3D]4f$	$^4F_{9/2}$	40.9155	9.802-13
295	$4s^2 4p^3(^2P)4d[^1F]4f$	$^2D_{3/2}$	40.9885	1.277-12
296	$4s^2 4p^3(^2P)4d[^3D]4f$	$^4F_{5/2}$	41.0028	1.109-12
297	$4s4p^5(^3P)4f$	$^4D_{1/2}$	41.0377	1.064-12
298	$4s^2 4p^3(^2P)4d[^3D]4f$	$^4G_{11/2}$	41.0404	1.391-12
299	$4s^2 4p^3(^2P)4d[^1F]4f$	$^2F_{7/2}$	41.0763	1.522-12
300	$4s4p^5(^3P)4f$	$^4D_{3/2}$	41.1978	9.994-13

Index	Configuration	Level	Energy	τ (s)
301	$4s^2 4p^3(^2P)4d[^1F]4f$	$^2P_{1/2}$	41.2134	1.027-12
302	$4s4p^5(^1P)4f$	$^2D_{3/2}$	41.2687	1.516-12
303	$4s^2 4p^3(^2P)4d[^1P]4f$	$^2G_{9/2}$	41.2760	2.039-12
304	$4s^2 4p^3(^2D)4d[^3G]4f$	$^2K_{13/2}$	41.2769	1.355-12
305	$4s^2 4p^3(^4S)4d[^5D]4f$	$^4P_{1/2}$	41.3689	2.310-12
306	$4s^2 4p^3(^2P)4d^2(^3P)$	$^2P^o_{1/2}$	41.3253	3.110-13
307	$4s4p^5(^3P)4f$	$^4D_{5/2}$	41.4078	1.508-12
308	$4s4p^5(^3P)4f$	$^4G_{7/2}$	41.4607	1.299-12
309	$4s4p^5(^3P)4f$	$^4F_{3/2}$	41.4655	1.604-12
310	$4s4p^5(^3P)4f$	$^4G_{9/2}$	41.4643	1.261-12
311	$4s4p^5(^3P)4f$	$^4G_{11/2}$	41.4665	1.148-12
312	$4s4p^5(^3P)4f$	$^2F_{5/2}$	41.4864	1.159-12
313	$4s^2 4p^2(^3P)4d^3(^2D)$	$^4D_{1/2}$	41.5531	8.096-13
314	$4s^2 4p^3(^2P)4d[^1F]4f$	$^2D_{5/2}$	41.6322	1.542-12
315	$4s4p^5(^3P)4f$	$^2F_{7/2}$	41.7308	1.194-12
316	$4s^2 4p^3(^2P)4d[^1F]4f$	$^2S_{1/2}$	41.7401	1.297-12
317	$4s4p^5(^3P)4f$	$^4D_{7/2}$	41.7288	1.108-12
318	$4s^2 4p^3(^2P)4d^2(^3P)$	$^2P^o_{3/2}$	41.6480	7.838-13
319	$4s^2 4p^3(^2P)4d[^3D]4f$	$^2H_{11/2}$	41.7475	8.543-13
320	$4s4p^5(^3P)4f$	$^2D_{3/2}$	41.7663	9.165-13
321	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6F_{3/2}$	41.8170	7.889-13
322	$4s4p^5(^3P)4f$	$^4G_{5/2}$	41.8470	1.226-12
323	$4s4p^5(^3P)4f$	$^2G_{9/2}$	41.9010	1.124-12
324	$4s^2 4p^3(^2P)4d^2(^3F)$	$^2D^o_{5/2}$	41.8290	6.023-13
325	$4s^2 4p^4(^1S)4f$	$^2F^o_{7/2}$	41.8711	8.508-13
326	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6F_{5/2}$	41.9804	7.472-13
327	$4s4p^5(^3P)4f$	$^2G_{7/2}$	42.0593	9.701-13
328	$4s^2 4p^2(^1D)4d^3(^4F)$	$^4H_{7/2}$	42.1444	9.123-13
329	$4s^2 4p^3(^2P)4d[^1P]4f$	$^2D_{3/2}$	42.1586	1.171-12
330	$4s^2 4p^3(^2P)4d^2(^1G)$	$^2F^o_{5/2}$	42.1726	3.686-13
331	$4s4p^5(^3P)4f$	$^2D_{5/2}$	42.3479	1.142-12
332	$4s4p^5(^3P)4f$	$^4F_{9/2}$	42.4077	1.057-12
333	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6G_{7/2}$	42.4498	1.401-12
334	$4s^2 4p^2(^3P)4d^3(^2P)$	$^4D_{5/2}$	42.4592	1.403-12
335	$4s^2 4p^2(^3P)4d^3(^4P)$	$^6P_{3/2}$	42.4851	1.642-12
336	$4s4p^5(^1P)4f$	$^2G_{7/2}$	42.5133	1.034-12
337	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6G_{9/2}$	42.5342	1.509-12
338	$4s4p^5(^1P)4f$	$^2F_{5/2}$	42.5732	1.029-12
339	$4s^2 4p^3(^4S)4d[^3D]4f$	$^2P_{3/2}$	42.6916	1.033-12
340	$4s4p^5(^1P)4f$	$^2G_{9/2}$	42.7559	1.061-12
341	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6F_{1/2}$	42.7647	9.289-13
342	$4s4p^5(^1P)4f$	$^2D_{5/2}$	42.7749	1.369-12
343	$4s^2 4p^2(^3P)4d^3(^4F)$	$^4F_{3/2}$	42.7873	1.075-12
344	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6G_{11/2}$	42.8022	1.796-12
345	$4s4p^5(^1P)4f$	$^2F_{7/2}$	42.8457	1.213-12
346	$4s^2 4p^2(^3P)4d^3(^4F)$	$^4F_{5/2}$	42.9412	7.338-13
347	$4s^2 4p^2(^1D)4d^3(^4F)$	$^4D_{1/2}$	42.9632	5.222-13
348	$4s^2 4p^2(^3P)4d^3(^2G)$	$^4G_{7/2}$	43.0064	1.172-12
349	$4s^2 4p^2(^1D)4d^3(^2H)$	$^2H_{9/2}$	43.0361	1.436-12
350	$4s^2 4p^2(^1D)4d^3(^2G)$	$^2I_{13/2}$	43.0840	1.570-12

Index	Configuration	Level	Energy	τ (s)
351	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6D_{3/2}$	43.1717	4.522-13
352	$4s^2 4p^2(^3P)4d^3(^4P)$	$^6D_{5/2}$	43.2639	4.831-13
353	$4s^2 4p^2(^3P)4d^3(^2F)$	$^4D_{7/2}$	43.3970	1.437-12
354	$4s^2 4p^2(^1D)4d^3(^2G)$	$^2H_{9/2}$	43.4078	1.532-12
355	$4s^2 4p^2(^3P)4d^3(^2H)$	$^4I_{11/2}$	43.4029	1.581-12
356	$4p^6 4d$	$^2D_{3/2}$	43.5244	5.304-13
357	$4s^2 4p^2(^3P)4d^3(^4P)$	$^6D_{7/2}$	43.4998	1.208-12
358	$4s^2 4p^2(^3P)4d^3(^4F)$	$^4G_{5/2}$	43.5981	1.021-12
359	$4s^2 4p^2(^3P)4d^3(^4F)$	$^6F_{9/2}$	43.6047	2.241-12
360	$4s^2 4p^3(^2D)4d[^3D]4f$	$^4G_{5/2}$	43.6906	2.902-12